



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2016

Malassezia Dermatitis - How do I manage this?

Rostaher, A

Posted at the Zurich Open Repository and Archive, University of Zurich
ZORA URL: <https://doi.org/10.5167/uzh-126301>
Conference or Workshop Item
Published Version

Originally published at:

Rostaher, A (2016). Malassezia Dermatitis - How do I manage this? In: 8th World Congress of Veterinary Dermatology, Bordeaux, France, 31 May 2016 - 4 June 2016.

MALASSEZIA DERMATITIS - HOW DO I MANAGE THIS

Ana Rostaher

Dermatology unit, Clinic for Small Animal Internal Medicine, Vetsuisse Faculty University of Zurich, Zurich, Switzerland

1. Introduction

Malassezia dermatitis is common in dogs, and less common in cats.^{1,2} *Malassezia* spp. are commensal yeasts that usually colonize the superficial epidermis of the skin and the ears. The non-lipid dependend *M. pachydermatis* and the lipid dependent *M. sympodialis*, *M. globosa*, *M. nana*, *M. slooffiae*, *M. restricta* and *M. furfur* are usually found on the skin or mucosae of dogs and cats.¹⁻³ Because *Malassezia* sp. do not invade subcorneally, it is hypothesized that the dermatitis results from inflammatory and/or hypersensitivity reactions to yeast enzymatic products and antigens.⁴ *Malassezia*-specific IgG and IgE seem to be higher in atopic then normal dogs pointing toward a possible involvement in the pathogenesis of atopic dermatitis.^{5,6} This was also shown in humans.⁷

Predisposing or triggering factors for *Malassezia* dermatitis development are as follows: increased humidity, presence of skin folds, endocrinopathies (hypothyroidism, diabetes mellitus, hyperadrenocorticism), increased populations of staphylococci, immunologic alterations (allergies, FIV, neoplasia, glucocorticoid therapy), and genetic predisposition (Basset hound, West Highland White Terrier, American Cocker Spaniel, and many other dog breeds, short-hair cat breeds such as Sphynx and Rex cats).^{1-3,8-12}

2. Clinical signs

Predominant skin lesion in dogs are erythema, scaling, greasiness, crusts, in chronic cases also lichenification and hyperpigmentation (localized or generalized).² Sometimes the disease is associated with papules and even furunculosis.² If affected, the claws show red-brown discoloration. Predilection site are lips, ear canals, groin, ventral neck, medial thighs, perianal area, interdigital skin and skin folds. In addition to skin changes, nearly all patients experience pruritus.² In the cat, *Malassezia* dermatitis should be on the differential list, in following clinical presentations:

- Otitis externa (often with accumulations of brown/black and waxy material)
- Chin acne
- Facial dermatitis
- Paronychia and/or red-brown discoloration of the claws
- Generalized erythematous scaly to waxy dermatitis^{2,8,11,13,14}

3. Diagnosis

Main differential diagnoses include allergic dermatitis, contact dermatitis, superficial staphylococcal folliculitis, demodicosis, dermatophytosis, scabies, cheyletiellosis, feline acne and epitheliotropic lymphoma. Remember that most of the dogs and cats with *Malassezia* dermatitis have concurrent dermatoses or a systemic disease, which should be screened for.

The diagnosis is made by cytological examination using direct impressions with a microscopic slide, swab, superficial scraping or cellophane tape strippings (for less accessible areas such as interdigital skin or dry lesions).² Whichever method is used, the glass slides are usually stained with Romanowsky stain (Diff-Quik®). Optionally, the samples (not scotch tape) can be heat

fixed before staining.¹⁵ This may prevent loss of some material and consequently important information during the staining procedure. If one is searching only infectious agents such as bacteria and *Malassezia* just the basophilic staining component may be used. Additional utilization of the eosinophilic stain reveals important information about the inflammatory response (Example eosinophils). The yeasts are best visualized at 40x or 100x magnification and are round to oval shaped with or without budding (classical “peanut shape”). The yeast diameter maybe from 3-8 um in diameter.² There is no definite yeast number to diagnose *Malassezia* dermatitis. Therefore, the diagnostic criteria for *Malassezia* dermatitis should always be interpreted in the clinical context. Some cases even require a therapeutical trial, to confirm the diagnosis. The fungal culture is of **no diagnostic value** in superficial infections, as dogs and cats can be healthy carriers. Important is to have typical clinical lesions with associated *Malassezia* organisms found on the cytological examination.

4. Treatment

Treatment options depend on the severity of lesions and the compliance of the owner/patient. Small focal areas can be treated with antifungal creams, wipes or ear drops. In multifocal or even generalized disease the topical therapy includes application of shampoos and rinses in mono- or combined preparations: chlorhexidin, ketoconazole, miconazole, climbazole, selenium sulfide, enilconazole, lime sulfur, acetic and boric acid. When topical therapy is impractical or ineffective, oral antifungal treatment may be instituted. The most commonly used drugs are itraconazole, ketoconazole, terbinafine.^{13,16-21} As the therapeutic levels of systemic antifungals persist in the skin for several days or weeks, pulse regimens are possible. Newer treatment modalities, such as killer peptides, are being described recently.²² Very important is to consider possible drug interactions when creating a multimodal therapeutic regimen (example allergic dogs treated with cyclosporine). As infections with *Malassezia* sp. often co-exist with staphylococcal infections, do not forget to diagnose and treat also these. Patients with a clinically relevant inflammatory component (usually allergic dogs), require additional anti-inflammatory drugs, such as glucocorticoids. Antifungal susceptibility testing is not routinely performed, but should be considered in refractory cases, as azole-resistant cases have been already encountered in veterinary medicine.²³

Dramatic clinical improvement is seen within 1 to 2 weeks (faster if topical and systemic medications are combined). The duration of the therapy depends on the resolution of clinical signs and should be continued 7-10 days beyond clinical cure, in average altogether 4 weeks.² Recurrent cases of *Malassezia* dermatitis are not unusual, as generally an underlying chronic disease is present. Frequent recurrences may require a maintenance protocol with utilization of either regular topical (once to twice weekly) or systemic antifungals (1-3 days/week pulses).

Again, to achieve a good clinical response to treatment, all efforts should be made to identify and correct the predisposing factors (allergy, hormonal disturbances, neoplasia, immune-deficiencies, etc).

References

1. Guillot J, Bond R. *Malassezia pachydermatis*: a review. *Med Mycol* 1999; 37:295-306.
2. Scott D, Miller W, Griffin C. Fungal and Algal Skin Diseases In: Scott DW, Miller W, Griffin CE, ed. *Muller and Kirk's Small Animal Dermatology* 7th ed. Missouri, USA: Elsevier, 2013 223-283.
3. Bond R. Superficial veterinary mycoses. *Clin Dermatol* 2010; 28:226-236.

4. Velegraki A, Cafarchia C, Gaitanis G et al. Malassezia infections in humans and animals: pathophysiology, detection, and treatment. *PLoS Pathog* 2015; 11:e1004523.
5. Chen TA, Halliwell RE, Pemberton AD et al. Identification of major allergens of *Malassezia pachydermatis* in dogs with atopic dermatitis and *Malassezia* overgrowth. *Vet Dermatol* 2002; 13:141-150.
6. Nuttall TJ, Halliwell RE. Serum antibodies to *Malassezia* yeasts in canine atopic dermatitis. *Vet Dermatol* 2001; 12:327-332.
7. Glatz M, Bosshard PP, Hoetzenecker W et al. The Role of *Malassezia* spp. in Atopic Dermatitis. *J Clin Med* 2015; 4:1217-1228.
8. Colombo S, Nardoni S, Cornegiani L et al. Prevalence of *Malassezia* spp. yeasts in feline nail folds: a cytological and mycological study. *Vet Dermatol* 2007; 18:278-283.
9. Forster-Van Hijfte MA, Curtis CF, White RN. Resolution of exfoliative dermatitis and *Malassezia pachydermatis* overgrowth in a cat after surgical thymoma resection. *J Small Anim Pract* 1997; 38:451-454.
10. Godfrey DR. A case of feline paraneoplastic alopecia with secondary *Malassezia*-associated dermatitis. *J Small Anim Pract* 1998; 39:394-396.
11. Ordeix L, Galeotti F, Scarpella F et al. *Malassezia* spp. overgrowth in allergic cats. *Vet Dermatol* 2007; 18:316-323.
12. Plant JD, Rosenkrantz WS, Griffin CE. Factors associated with and prevalence of high *Malassezia pachydermatis* numbers on dog skin. *J Am Vet Med Assoc* 1992; 201:879-882.
13. Bensignor E. Treatment of *Malassezia* overgrowth with itraconazole in 15 cats. *Vet Rec* 2010; 167:1011-1012.
14. Nardoni S, Mancianti F, Rum A et al. Isolation of *Malassezia* species from healthy cats and cats with otitis. *J Feline Med Surg* 2005; 7:141-145.
15. Griffin JS, Scott DW, Erb HN. *Malassezia* otitis externa in the dog: the effect of heat-fixing otic exudate for cytological analysis. *J Vet Med A Physiol Pathol Clin Med* 2007; 54:424-427.
16. Ahman S, Perrins N, Bond R. Treatment of *Malassezia pachydermatis*-associated seborrhoeic dermatitis in Devon Rex cats with itraconazole--a pilot study. *Vet Dermatol* 2007; 18:171-174.
17. Cavana P, Peano A, Petit JY et al. A pilot study of the efficacy of wipes containing chlorhexidine 0.3%, climbazole 0.5% and Tris-EDTA to reduce *Malassezia pachydermatis* populations on canine skin. *Vet Dermatol* 2015; 26:278-e261.
18. Cavana P, Petit JY, Perrot S et al. Efficacy of a 2% climbazole shampoo for reducing *Malassezia* population sizes on the skin of naturally infected dogs. *J Mycol Med* 2015; 25:268-273.
19. Gimmler JR, White AG, Kennis RA et al. Determining canine skin concentrations of terbinafine to guide the treatment of *Malassezia* dermatitis. *Vet Dermatol* 2015; 26:411-416, e495-416.
20. Negre A, Bensignor E, Guillot J. Evidence-based veterinary dermatology: a systematic review of interventions for *Malassezia* dermatitis in dogs. *Vet Dermatol* 2009; 20:1-12.
21. Peano A, Gallo MG. Management of *Malassezia*-related diseases in the dog. *Parassitologia* 2008; 50:85-88.
22. Cafarchia C, Immediato D, Paola GD et al. In vitro and in vivo activity of a killer peptide against *Malassezia pachydermatis* causing otitis in dogs. *Med Mycol* 2014; 52:350-355.
23. Nijima M, Kano R, Nagata M et al. An azole-resistant isolate of *Malassezia pachydermatis*. *Vet Microbiol* 2011; 149:288-290.